



Frederick P. Fish
1855-1930

W.K. Richardson
1859-1951

11-28-00
FISH & RICHARDSON P.C., P.A.

November 27, 2000

Attorney Docket No.: 07039-170002

Box Patent Application

Commissioner for Patents
Washington, DC 20231

Presented for filing is a new continuation patent application of:

Applicant: JORG J. GORONZY AND CORNELIA M. WEYAND

Title: METHODS AND MATERIALS FOR EVALUATING RHEUMATOID
ARTHRITIS

The prior application is assigned of record to Mayo Foundation For Medical
Education and Research, a Minnesota corporation, by virtue of an assignment
submitted to the Patent and Trademark Office and recorded on November 30,
1999 at 010421/0339.

Enclosed are the following papers, including those required to receive a filing date
under 37 CFR §1.53(b):

	<u>Pages</u>
Specification	45
Claims	8
Abstract	1
Declaration	2
Drawing(s)	6

Enclosures:

- Preliminary amendment, 4 pages.
- Small entity statement. This application is entitled to small entity status.

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No EL 427178087US

I hereby certify under 37 CFR §1.10 that this correspondence is being
deposited with the United States Postal Service as Express Mail Post
Office to Addressee with sufficient postage on the date indicated below
and is addressed to the Commissioner for Patents, Washington,
D C 20231.

November 27, 2000
Date of Deposit

Signature

Vince Defante

Typed or Printed Name of Person Signing Certificate

3300 Dain Rauscher Plaza
60 South Sixth Street
Minneapolis, Minnesota
55402

Telephone
612 335-7070

Facsimile
612 288-5500

Web Site
www.fr.com



BOSTON

DALLAS

DELAWARE

NEW YORK

SAN DIEGO

SILICON VALLEY

TWIN CITIES

WASHINGTON, DC

09/23/00 14:27:00

— Postcard.

Basic filing fee	\$355
Total claims in excess of 20 times \$9	\$0
Independent claims in excess of 3 times \$40	\$0
Fee for multiple dependent claims	\$0
Total filing fee:	\$355

If this application is found to be incomplete, or if a telephone conference would otherwise be helpful, please call the undersigned at (612) 335-5070.

Kindly acknowledge receipt of this application by returning the enclosed postcard.

Please send all correspondence to:

MARK S. ELLINGER, PH.D.
Fish & Richardson P.C., P.A.
60 South Sixth Street
Suite 3300
Minneapolis, MN 55402

Respectfully submitted,

Q. R. A. F. - III

J. Patrick Finn III, Ph.D.
Reg. No. 44,109

Enclosures
JXF/jyh
60027213 doc

Applicant or Patentee: Jorg J. Goronzy

Serial or Patent No.: 09/387,467

Filed or Issued: September 1, 1999

For: METHODS AND MATERIALS FOR EVALUATING RHEUMATOID ARTHRITIS

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
(37 CFR 1.9(f) and 1.27(d)) — NONPROFIT ORGANIZATION**

I hereby declare that I am an official empowered to act on behalf of the nonprofit organization identified below:

Name of Organization: Mayo Foundation for Medical Education and Research

Address of Organization: 200 First Street S.W.
Rochester, Minnesota 55905

Type of Organization:

- ☐ University or Other Institution of Higher Education
- ☒ Tax Exempt Under Internal Revenue Service Code (26 SC 501(a) and 501(c)(3))
- ☐ Nonprofit Scientific or Educational Under Statute of State of the United States of America
(Name of State:) (Citation of Statute:)
- ☐ Would qualify as tax exempt under Internal Revenue Service Code (26 USC 501(a) and 501(c)(3)) if located in the United States of America
- ☐ Would qualify as nonprofit scientific or educational under Statute of State of the United States of America if located in the United States of America
(Name of State:) (Citation of Statute:)

I hereby declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR 1.9(e) for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code with regard to the invention entitled METHODS AND MATERIALS FOR EVALUATING RHEUMATOID ARTHRITIS by inventor(s) Jorg J. Goronzy and Cornelia M. Weyand described in:

- ☐ the specification filed herewith.
- ☒ application serial no. 09/387,467, filed September 1, 1999.
- ☐ patent no. _____, issued _____.

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization with regard to the above identified invention.

If the rights held by the nonprofit organization are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR 1.9(c) or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

*NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

Full Name: _____

Address: _____

☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status when any new rule 53 application is filed or prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

Name:	Rick F. Colvin
Title:	Assistant Treasurer
Address:	200 First Street S.W. Rochester, Minnesota 55905
Signature:	<i>Rick F. Colvin</i> Date: 11/5/99

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant : Jorg J. Goronzy et al. Art Unit : Unknown
Serial No. : To Be Assigned Examiner : Unknown
Filed : Concurrently herewith
Continuation of
Serial No. : 09/387,467
Title : METHODS AND MATERIALS FOR EVALUATING RHEUMATOID
ARTHRITIS

Commissioner for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Prior to examination, please amend the application as follows:

In the Specification:

On page 1, line 6, after "This application"; please insert --is a continuation application of U.S. Patent Application Serial Number 09/387,467, filed September 1, 1999, which--.

In the Claims:

Please cancel claims 1-47 without prejudice.

Please add new claims 48-62 as follows.

--48. A method for determining the predisposition of a rheumatoid arthritis patient to develop severe disease, said method comprising:

a) comparing the frequency of CD4⁺/CD28^{null} cells in said patient to a reference frequency to obtain information about said rheumatoid arthritis condition, and

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No EL 427178087US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

November 27, 2000

Date of Deposit

Signature

Vince Defante

Typed or Printed Name of Person Signing Certificate

00427178087US

b) determining if said patient is predisposed to develop severe disease based on said information and the presence or absence of a polymorphism in an HLA-DRB1 allele in said patient.

49. The method of claim 48, wherein said frequency of CD4⁺/CD28^{null} cells comprises the percent of CD4⁺ cells that are CD28 negative.

50. The method of claim 48, wherein said reference frequency is derived from the CD4⁺/CD28^{null} cell frequency from a population.

51. The method of claim 50, wherein said population comprises a population of patients having a diffuse rheumatoid arthritis condition.

52. The method of claim 50, wherein said population comprises a population of patients having a follicular rheumatoid arthritis condition.

53. The method of claim 50, wherein said population comprises a population of patients having a granulomatous rheumatoid arthritis condition.

54. The method of claim 50, wherein said population comprises a population of healthy individuals.

55. The method of claim 50, wherein said population comprises a population of patients having subcutaneous nodules.

56. The method of claim 50, wherein said population comprises a population of patients having extra-articular involvement.

57. The method of claim 50, wherein said population comprises a population of patients having major joint destruction.

CONFIDENTIAL

58. The method of claim 48, wherein said polymorphism comprises an HLA-DRB1 allele that encodes a polypeptide having an uncharged amino acid at position 74.

59. The method of claim 48, wherein said polymorphism comprises an HLA-DRB1 allele that encodes a polypeptide free from negatively charged amino acids at positions 70 and 71.

60. A method for determining the predisposition of a rheumatoid arthritis patient to develop severe disease, said method comprising:

- a) determining the frequency of $CD4^+/CD28^{null}$ cells in said patient,
- b) determining the presence or absence of a polymorphism in an HLA-DRB1 allele in said patient,
- c) comparing said frequency of $CD4^+/CD28^{null}$ cells to a reference frequency to obtain information about said rheumatoid arthritis condition, and
- d) determining if said patient is predisposed to develop severe disease based on said information and said presence or absence of said polymorphism.

61. A kit for determining the predisposition of a rheumatoid arthritis patient to develop severe disease, said kit comprising:

- a) a first binding pair member, wherein said first binding pair member has specific binding affinity for a $CD4^+/CD28^{null}$ cell marker such that the frequency of $CD4^+/CD28^{null}$ cells in said patient is determinable, and
- b) an oligonucleotide primer, wherein said oligonucleotide primer has specific binding affinity for at least a portion of the locus containing an HLA-DRB1 allele such that the a polymorphism of said HLA-DRB1 allele in said patient is determinable.

62. The kit of claim 61, wherein said kit comprises a reference chart, wherein said reference chart contains information about $CD4^+/CD28^{null}$ cell frequencies such that said predisposition is determinable based on said frequency of $CD4^+/CD28^{null}$ cells in said patient and said polymorphism of said HLA-DRB1 allele.--

REMARKS


Claims 1-47 have been cancelled herein without prejudice. In addition, claims 48-62 have been added herein. Thus, claims 48-62 are pending. The specification as filed provides support for these amendments. For example, page 4, lines 4-13 disclose that a patient's HLA-DRB1 alleles and frequency of CD4⁺/CD28^{null} cells can be used to determine the predisposition to develop severe disease. Thus, no new matter has been added.

Applicants submit that claims 48-62 are in condition for examination, which action is requested. Please apply any charges or credits to Deposit Account No. 06-1050.

Respectfully submitted,

Date:

November, 27, 2000



J. Patrick Finn III, Ph.D.
Reg. No. 44,109

Fish & Richardson P.C., P.A.
60 South Sixth Street
Suite 3300
Minneapolis, MN 55402
Telephone: (612) 335-5070
Facsimile: (612) 288-9696

60026974 doc

002277 0002260

[illegible][illegible]

Date of Deposit September 1, 1999
I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office To Addressee" with sufficient postage on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231

Date of Deposit September 1, 1999
I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office To Addressee" with sufficient postage on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231

B. William
Baker

**METHODS AND MATERIALS FOR EVALUATING
RHEUMATOID ARTHRITIS**

RELATED APPLICATIONS

This application claims priority from U.S. Provisional Application Serial No.
60/098,718, filed September 1, 1998.

BACKGROUND

1. Technical Field

The invention relates to methods and materials for evaluating rheumatoid arthritis as well as for determining an individual's predisposition to have severe rheumatoid arthritis disease.

2. Background Information

Rheumatoid arthritis (RA) affects individuals in the prime of their life and is feared because of its potential to cause chronic pain and irreversible damage of tendons, ligaments, joints, and bones. The symmetrical involvement of small peripheral joints has an enormous impact on hand and foot functions and poses therapeutic challenges that cannot be easily overcome by joint replacement. Also, systemic manifestations of RA are not rare and can range from relatively minor problems, such as rheumatoid nodules, to life-threatening organ disease.

In addition, RA is a systemic inflammatory disease that primarily manifests itself as synovial inflammation of diarthrodial joints. The typical histopathological changes include dense infiltration of the synovial membrane by mononuclear cells, neoangiogenesis, and hypertrophy and hyperplasia of the synovial lining (Harris ED (ed); Rheumatoid Arthritis, Philadelphia, WB Saunders Co., pp.3-212 (1997); and Hale LP, Haynes BF: Pathology of rheumatoid arthritis and associated disorders. Arthritis and Allied Conditions. A textbook of Rheumatology. Edited by Koopman WJ. Baltimore, Williams & Wilkins, pp.993-1016 (1997)). The etiopathogenesis of the syndrome is not understood. Several lines of evidence

support a central role of T lymphocytes in the disease-specific pathogenic events (Todd, *et al. Science*, 240:1003-1009 (1988); Panayi, *et al., Arthritis Rheum*, 35:729-735 (1992); and Goronzy JJ, Weyand CM: *Rheum Dis Clin North Am*, 21:655-674 (1995)). An alternative hypothesis, namely, that macrophages are the pivotal cell type in rheumatoid synovitis, has also been proposed (Firestein GS, Zvaifler NJ: *Arthritis Rheum* 33:768-773 (1990); and Burmester, *et al., Arthritis Rheum*, 40:5-18 (1997)). Whether only T cells or only macrophages or both are the causative elements in RA remains a matter of controversy (Feldmann, *et al., Cell*, 85:307-310 (1996); and Fox, *Arthritis Rheum* 40:598-609 (1997)).

RA is primarily a clinical diagnosis. Symmetrical joint involvement, dominant manifestations in peripheral joints, rheumatoid factor production, and the formation of rheumatoid nodules are considered when the diagnosis is made (Arnett, *et al., Arthritis Rheum*. 31:315-324 (1988)). The histological appearance of the synovium varies quite extensively and the pathological findings are usually not helpful in distinguishing RA from other inflammatory arthropathies (Hale LP, Haynes BF: Pathology of rheumatoid arthritis and associated disorders. *Arthritis and Allied Conditions. A textbook of Rheumatology*. Edited by Koopman WJ. Baltimore, Williams & Wilkins, pp.993-1016 (1997)). In addition, no information is available on the mechanisms underlying the topographical arrangement of the inflammatory infiltrate in the rheumatoid synovium.

Therapeutic management of RA has steadily improved over the last decades, mostly due to the recognition that destruction caused by chronic inflammation is irreversible and that only early and aggressive intervention can enhance therapeutic benefit. Consequently, RA patients are now being treated early in the disease course and disease modifying agents are widely used. Despite these successes, major challenges remain. Presently, no curative intervention is available, side effects of therapies are significant, and the disease may still progress while the patient is being treated.

SUMMARY

The invention involves methods and materials for evaluating rheumatoid arthritis in a patient. Specifically, the invention provides methods and materials for classifying a rheumatoid arthritis condition as diffuse, follicular, or granulomatous. In addition, the

invention provides methods and materials for determining if an individual suffering from a rheumatoid arthritis condition will develop severe disease. Useful indicators for severe disease include, without limitation, subcutaneous nodule formation (nodularity) and extra-articular involvement.

5 The invention is based on the discovery that the level of particular cytokines within tissue or the histological appearance of tissue or both can be used to classify a rheumatoid arthritis condition as diffuse, follicular, or granulomatous. This classification is important since granulomatous patients are more susceptible to severe rheumatoid arthritis disease. Severe rheumatoid arthritis disease can involve, without limitation, major organ involvement,
10 which can be life threatening, and major joint destruction, which can be crippling. Thus, proper classification of a granulomatous rheumatoid arthritis condition can help provide clinicians and patients with information that can be used to determine adequate treatments.

 Specifically, the invention involves analyzing a synovial tissue biopsy for particular cytokines such as IL-4, IL-10, and IFN- γ , or for particular histological characteristics, or
15 both. For example, a patient can be classified as having a diffuse condition when tissue from that patient contains low levels of IL-4, IL-10, and IFN- γ , or as having a follicular condition when tissue from that patient contains high levels of IL-10 and IFN- γ and low levels of IL-4. A particular level of a particular cytokine can be determined to be high or low based on the levels measured from various populations. Such populations can include, without limitation,
20 populations of patients with a diffuse condition, follicular condition, or granulomatous condition, patients with subcutaneous nodule formation, patients with extra-articular involvement, patients with major joint destruction, and healthy individuals.

 The invention also is based on the discovery that patients presenting similar rheumatoid arthritis symptoms can have different levels of particular cytokines within their
25 tissue or different histological appearance of their tissue. Thus, determining the tissue cytokine profile or the histological characteristics of a synovial tissue sample from a patient can be used to determine the proper treatment protocol. For example, two patients having similar rheumatoid arthritis symptoms may have different levels of IL-10 within their tissue. The patient with low levels may benefit from a treatment of IL-10 while the patient with high
30 levels of IL-10 may benefit from treatment with IL-10 inhibitors such as anti-IL-10

antibodies. Thus, determining the tissue cytokine profile or the histological characteristics of a tissue sample from a patient can help provide clinicians and patients with information that can be used to determine adequate treatments.

In addition, the invention is based on the discovery that the predisposition to develop severe disease can be determined in patients classified as having diffuse or follicular disease by analyzing the patient's HLA-DRB1 alleles or frequency of CD4⁺/CD28^{null} cells or both. Determining a patient's predisposition to develop severe disease is important since it allows clinicians and patients to plan and treat accordingly. Again, severe rheumatoid arthritis disease can involve, without limitation, extra-articular involvement and major joint destruction. Moreover, analyzing a patient's HLA-DRB1 alleles and/or frequency of CD4⁺/CD28^{null} cells can be used to determine whether a patient classified as having diffuse or follicular disease with a propensity for severe disease will have major organ damage, major joint destruction, or both.

Specifically, the invention involves determining whether a patient having a diffuse or follicular condition has zero, one, or two HLA-DRB1 alleles that are associated with RA and/or whether that patient has an elevated frequency of CD4⁺/CD28^{null} cells. For example, HLA-DRB1 alleles that are associated with RA can be any allele having a polymorphism associated with rheumatoid arthritis including HLA-DRB1 alleles that encode polypeptides having an uncharged amino acid at position 74, no negatively charged amino acid at position 70, and a positively charged amino acid at position 71. Patients having RA-associated polymorphisms for one or both HLA-DRB1 alleles or having an elevated frequency of CD4⁺/CD28^{null} cells can be classified as having the potential to form severe disease.

In general, the invention features a method for diagnosing a rheumatoid arthritis condition in a patient. The method includes determining the level of a cytokine (e.g., IL-4, IL-10, and IFN- γ) within a sample from the patient, comparing the level of the cytokine to a reference level to obtain information about the rheumatoid arthritis condition, and classifying the rheumatoid arthritis condition as a diffuse, follicular, or granulomatous condition based on the information. The sample can be a tissue biopsy (e.g., a synovial tissue biopsy). The reference level can be the median level of the cytokine found in tissue samples derived from a population. The population can include a population of patients having a diffuse rheumatoid

arthritis condition, a population of patients having a follicular rheumatoid arthritis condition, a population of patients having a granulomatous rheumatoid arthritis condition, a population of healthy individuals, a population of patients having subcutaneous nodules, a population of patients having extra-articular involvement, or a population of patients having major joint destruction.

In another embodiment, the invention features a method for determining the predisposition of a rheumatoid arthritis patient to develop severe disease. The method includes determining the level of a cytokine (e.g., IL-4, IL-10, and IFN- γ) within a sample from the patient, determining the frequency of CD4⁺/CD28^{null} cells in the patient, comparing the level of the cytokine to a reference level and the frequency of CD4⁺/CD28^{null} cells to a reference frequency to obtain information about the predisposition, and determining if the patient is predisposed to develop severe disease based on the information. The sample can be a tissue biopsy (e.g., a synovial tissue biopsy). The reference level can be the median level of the cytokine found in tissue samples derived from a population. The population can include a population of patients having a diffuse rheumatoid arthritis condition, a population of patients having a follicular rheumatoid arthritis condition, a population of patients having a granulomatous rheumatoid arthritis condition, a population of healthy individuals, a population of patients having subcutaneous nodules, a population of patients having extra-articular involvement, or a population of patients having major joint destruction. The frequency of CD4⁺/CD28^{null} cells can be the percent of CD4⁺ cells that are CD28 negative. In addition, the reference frequency can be derived from the CD4⁺/CD28^{null} cell frequency from a population.

Another embodiment of the invention features a method for determining the predisposition of a rheumatoid arthritis patient to develop severe disease. The method includes determining the level of a cytokine (e.g., IL-4, IL-10, and IFN- γ) within a sample from the patient, comparing the level of the cytokine to a reference level to obtain information about the rheumatoid arthritis condition, determining the presence of a polymorphism in an HLA-DRB1 allele in the patient, and determining if the patient is predisposed to develop severe disease based on the information and the presence of the polymorphism. The sample can be a tissue biopsy (e.g., a synovial tissue biopsy). The reference level can be the median

level of the cytokine found in tissue samples derived from a population. The population can include a population of patients having a diffuse rheumatoid arthritis condition, a population of patients having a follicular rheumatoid arthritis condition, a population of patients having a granulomatous rheumatoid arthritis condition, a population of healthy individuals, a population of patients having subcutaneous nodules, a population of patients having extra-articular involvement, or a population of patients having major joint destruction. The polymorphism can include an HLA-DRB1 allele that encodes a polypeptide having an uncharged amino acid at position 74, or an HLA-DRB1 allele that encodes a polypeptide free from negatively charged amino acids at positions 70 and 71.

Another embodiment of the invention features a method for determining the predisposition of a rheumatoid arthritis patient to develop severe disease. The method includes determining the level of a cytokine within a sample from the patient, determining the frequency of CD4⁺/CD28^{null} cells in the patient, comparing the level of the cytokine to a reference level and the frequency of CD4⁺/CD28^{null} cells to a reference frequency to obtain information about the rheumatoid arthritis condition, determining the presence of a polymorphism in an HLA-DRB1 allele in the patient, and determining if the patient is predisposed to develop severe disease based on the information and the presence of the polymorphism.

In another aspect, the invention features a kit for providing diagnostic information about a rheumatoid arthritis condition in a patient. The kit contains a binding pair member and a reference chart. The binding pair member has specific binding affinity for a cytokine such that the level of the cytokine within a sample from the patient is determinable, and the reference chart contains information about cytokine levels such that an indication of the diffuse, follicular, or granulomatous nature of the rheumatoid arthritis condition is determinable based on the level of the cytokine within the sample.

In another embodiment, the invention features a kit for providing diagnostic information about a rheumatoid arthritis condition in a patient. The kit contains a binding pair member and a reference chart. The binding pair member has specific binding affinity for a nucleic acid sequence encoding a cytokine such that the level of the cytokine within a sample from the patient is determinable, and the reference chart contains information about

cytokine levels such that an indication of the diffuse, follicular, or granulomatous nature of the rheumatoid arthritis condition is determinable based on the level of the cytokine within the sample.

In another embodiment, the invention features a kit for determining the predisposition of a rheumatoid arthritis patient to develop severe disease. The kit contains a first binding pair member, a second binding pair member, and a reference chart. The first binding pair member has specific binding affinity for a cytokine or nucleic acid encoding the cytokine such that the level of the cytokine within a sample from the patient is determinable. The second binding pair member has specific binding affinity for a CD4⁺/CD28^{null} cell marker such that the frequency of CD4⁺/CD28^{null} cells in the patient is determinable. The reference chart contains information about cytokine levels and CD4⁺/CD28^{null} cell frequencies such that an indication of the predisposition is determinable based on the level of the cytokine within the sample and the frequency of CD4⁺/CD28^{null} cells in the patient.

In another embodiment, the invention features a kit for determining the predisposition of a rheumatoid arthritis patient to develop severe disease. The kit contains a binding pair member, an oligonucleotide primer, and a reference chart. The binding pair member has specific binding affinity for a cytokine or nucleic acid encoding the cytokine such that the level of the cytokine within a sample from the patient is determinable. The oligonucleotide primer has specific binding affinity for at least a portion of the locus containing an HLA-DRB1 allele such that a polymorphism of HLA-DRB1 allele in the patient is determinable. The reference chart contains information about cytokine levels such that an indication of the predisposition is determinable based on the level of the cytokine within the sample and the polymorphism of the patient. The kit can contain a plurality of the oligonucleotide primers.

In another embodiment, the invention features a kit for determining the predisposition of a rheumatoid arthritis patient to develop severe disease. The kit contains a first binding pair member, a second binding pair member, an oligonucleotide primer, and a reference chart. The first binding pair member has specific binding affinity for a cytokine or nucleic acid encoding the cytokine such that the level of the cytokine within a sample from the patient is determinable. The second binding pair member has specific binding affinity for a CD4⁺/CD28^{null} cell marker such that the frequency of CD4⁺/CD28^{null} cells in the patient is

determinable. The oligonucleotide primer has specific binding affinity for at least a portion of the locus containing an HLA-DRB1 allele such that the a polymorphism of the HLA-DRB1 allele in the patient is determinable. The reference chart contains information about cytokine levels and CD4⁺/CD28^{null} cell frequencies such that an indication of the predisposition is determinable based on the level of the cytokine within the sample, the frequency of CD4⁺/CD28^{null} cells in the patient, and the polymorphism of the HLA-DRB1 allele.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

DESCRIPTION OF DRAWINGS

Figure 1 contains two bar graphs plotting the number of cytokine transcripts (IFN- γ and IL-4) per β -actin transcripts for diffuse, follicular, and granulomatous synovial tissues.

Figure 2 contains two bar graphs plotting the number of cytokine transcripts (IL-1 β and TNF- α) per β -actin transcripts for diffuse, follicular, and granulomatous synovial tissues.

Figure 3 contains two bar graphs plotting the number of cytokine transcripts (IL-10 and TGF- β 1) per β -actin transcripts for diffuse, follicular, and granulomatous synovial tissues.

Figure 4 contains two graphs plotting CD28 expression against CD4 expression for a normal control individual (left, <1% of CD4⁺ T cells are CD28 negative) and a RA patient (right, about 9% of CD4⁺ T cells are CD28 negative).

Figure 5 is a bar graph depicting the correlation between the frequencies of CD4⁺/CD28^{null} T cells and extra-articular rheumatoid arthritis. The line corresponds to the

median, the box to the 25th and 75th percentiles, and the whiskers to the 10th and 90th percentiles.

Figure 6 is a chart depicting possible genetic factors of disease heterogeneity in rheumatoid arthritis.

5

DETAILED DESCRIPTION

The invention provides methods and materials for evaluating rheumatoid arthritis in a patient. Specifically, the invention provides methods and materials for classifying a rheumatoid arthritis condition as diffuse, follicular, or granulomatous. In addition, the invention provides methods and materials for determining a patient's predisposition to develop severe rheumatoid arthritis disease.

RA can be classified as diffuse, follicular, or granulomatous based on either the profile of cytokines produced within a synovial tissue sample or the histological characteristics of a synovial tissue sample. Any method can be used to quantify specific cytokine expression within synovial tissue including methods that measure cytokine mRNA or cytokine polypeptide. For example, PCR, competitive PCR, PCR-ELISA, and in situ hybridization techniques can be used to measure cytokine mRNA. In addition, ELISA, immunohistochemistry and other immuno assays can be used to measure cytokine polypeptide. Such cytokines can be, without limitation, IFN- γ , IL-2, IL-4, IL-5, IL-6, IL-10, and IL-12. The histological characteristics of a synovial tissue sample can be determined using common staining techniques that reveal the presence of macrophages and lymphocytes such as H & E staining and immunocytochemistry.

If the cytokine profiles or histological characteristics indicate that a patient has granulomatous disease then that patient also can be classified as being predisposed to develop a severe rheumatoid arthritis condition including major organ involvement and major joint destruction since granulomatous disease was discovered herein to correlate with severe disease. If the cytokine profiles or histological characteristics indicate that a patient has diffuse or follicular disease then further analysis can be performed to determine that patient's predisposition to develop severe disease. This further analysis can involve analyzing the HLA-DRB1 locus for alleles having polymorphisms associated with RA and/or determining

the CD4⁺/CD28^{null} T cell frequency.

Determining whether a patient has zero, one, or two HLA-DRB1 alleles that contain a polymorphism associated with RA can be accomplished using commonly known methods such as PCR, PCR-ELISA, and sequencing. For example, any method provided by the International Histocompatibility Workshop can be used as well as any commercially available methods of materials obtained from, for example, One Lambda (Los Angeles, CA) or Biotest Diagnostics (Denville, NJ). Such HLA-DRB1 alleles can include, without limitation, HLA-DRB1 alleles that encode a polypeptide having an uncharged amino acid at position 74, no negatively charged amino acid at position 70, and a positively charged amino acid at position 71.

Any method can be used to determine the frequency of CD4⁺/CD28^{null} T cells within a patient. For example, a binding pair member having specificity for a marker found on CD4⁺/CD28^{null} cells can be used to determine the frequency of those cells. Such a method can involve using the combination of two antibodies, one having specificity for CD4 and the other having specificity for CD28. For example, FACS can be used with CD4 and CD28 specific antibodies to determine the percent of CD4⁺ cells that are CD28 negative. Further, CD3 antibodies can be used to distinguish CD4⁺ T cells from CD4⁺ macrophages.

Once a frequency of CD4⁺/CD28^{null} cells in a patient is determined, that frequency can be compared to a reference frequency to obtain information about the patient's RA condition. Typically, a reference frequency is derived from the CD4⁺/CD28^{null} cell frequencies determined for a population of individuals. For example, a reference frequency can be a median percent of CD4⁺ cells that are CD28 negative as derived from a population. The population can be a population of healthy individuals or patients having subcutaneous nodularity, extra-articular involvement, major joint destruction, diffuse disease, follicular disease, and/or granulomatous disease. The information obtained by this comparison can allow the patient's RA condition to be classified as either severe or not severe. For example, if the frequency of CD4⁺/CD28^{null} cells from a patient corresponds to the reference frequency of CD4⁺/CD28^{null} cells derived from a population of healthy individuals and that patient has diffuse disease and no RA-associated HLA-DRB1 alleles, then that patient would not be classified as being predisposed to develop severe RA disease. Likewise, if the frequency of

CD4⁺/CD28^{null} cells from a patient corresponds to the reference frequency of CD4⁺/CD28^{null} cells derived from a population of patients having subcutaneous nodularity and that patient has diffuse and no RA-associated DRB1 alleles, then that patient can be classified as being predisposed to develop severe RA disease. Typically, a patient having diffuse RA disease with greater than about 2.0% of their CD4⁺ cells being CD28 negative has a predisposition to develop severe RA disease.

A reference chart is any chart that contains information about cytokine levels, HLA-DRB1 alleles, or CD4⁺/CD28^{null} T cell frequencies. For example, a reference chart can contain information about the average frequency of CD4⁺/CD28^{null} cells found in a particular population of individuals or patients. It will be appreciated that a reference chart can be presented or contained on any type of medium (e.g., paper or electronic formats). It is to be understood that an electronic format can be obtained via a software program or an accessible database site (e.g., an Internet site). For the purpose of this invention, kits containing a reference chart include kits having directions (e.g., access codes or Internet address information) for accessing reference charts in an electronic format.

A binding pair member is any molecule that specifically binds another molecule including, without limitation, antibodies, antibody fragments that have binding specificity, ligands, nucleic acid, receptors, lectins, chelating agents, ions, and the like.

The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

EXAMPLES

Example 1 - Three Distinct Histopathological Patterns in Rheumatoid Synovitis

The pattern of cytokines produced in the rheumatoid synovium was found to be predictive of the morphological appearance of the disease. Specifically, rheumatoid synovitis was found to be a heterogeneous entity with three distinct histologically defined phenotypes. The phenotypic heterogeneity correlated to a specific combination of T-cell- and macrophage-derived cytokines, raising the possibility that several pathomechanisms may cause an RA-like syndrome.

1. Study Population

Fresh synovial tissue was obtained from 21 consecutive patients with active RA who fulfilled the American College of Rheumatology 1987 revised criteria for RA (Arnett, *et al.*, *Arthritis Rheum*, pp.31:315-324 (1988)) and who underwent joint surgery.

2. Histopathological Evaluations

Hematoxylin and eosin sections of the tissue samples were analyzed for the organizational structure of the inflammatory infiltrate with particular attention to the topographical arrangement of T cells, B cells, and macrophages as well as the degree of angiogenesis and the relationship of the mononuclear infiltrate to the subsynovial lining. All tissue samples were reviewed by a pathologist who was unaware of any clinical, seriological, or immunohistological findings. To control for patchiness of the inflammation and intra-specimen variation, multiple independent specimens were included from ten patients with the pathologist blinded to the identity of the specimens. There was concordance for the identification of follicles and granulomatous lesions for all of the independently graded specimens.

3. Immunohistochemistry

Frozen tissue samples embedded in OCT (Miles, Elkhart, IN) were cut into 5- μ m sections, mounted on the gel-coated slides (Superfrost/Plus, Fisher Scientific, Pittsburgh, PA), and dried in a 37°C desiccator. Slides were stored at -70°C. Before staining, slides were fixed in acetone for ten minutes, air dried, and fixed in a 1% paraformaldehyde/EDTA, pH 7.2, for three minutes. Endogenous peroxidase was blocked with 0.3% H₂O₂ in 0.1% sodium azide. Nonspecific binding was blocked with 5% rabbit serum (Life Technologies, Grand Island, NY) for 15 minutes. Sections were stained with monoclonal mouse anti-interferon (IFN)- γ Ab, 1:100 (Genzyme Diagnostics, Cambridge, MA) for 60 minutes or monoclonal mouse anti-CD20 Ab, 1:40 (Dako, Carpinteria, CA) for 30 minutes at room temperature. After incubation with biotinylated rabbit anti-mouse antibody, 1:300 (Dako), the slides were developed with streptavidin-peroxidase, 1:250 (Dako) and 3,3'-diaminobenzidine tetrahydrochloride (Sigma Chemical Co., St. Louis, MO).

Slides stained with anti-IFN- γ were washed in 0.5% Triton X-100 in PBS for ten minutes. Nonspecific binding was blocked for 15 minutes with 5% normal goat serum (Life Technologies). Sections were stained with a monoclonal mouse anti-CD45RO Ab, 1:60 (Dako), for 30 minutes at room temperature. After incubation with a biotinylated rabbit anti-mouse antibody, 1:300 (Dako), the slides were developed with a Vectastain ABC-AP kit and alkaline phosphatase substrate kit 1 (Vector Laboratories, Burlingame, CA). Negative controls without primary Ab were processed in parallel. Sections were counterstained with hematoxylin and permanently mounted in Cytoseal (Stephens Scientific, Riverdale, NJ).

4. Cytokine Measurement

Total RNA was extracted from synovial tissue by using a commercially available kit (Trizol, Life Technologies). cDNA from synovial tissue specimens was adjusted to contain equal numbers of β -actin transcripts. Adjusted cDNA was amplified under nonsaturating conditions with cytokine-specific primers (Table 1) by polymerase chain reaction (PCR) in parallel with a standard with a known number of cytokine sequences as described (Weyand, *et al.*, *Arthritis Rheum*, pp.40:19-26 (1997)). Primers were chosen to amplify cDNA specifically and not genomic DNA. To achieve this goal, several primers had been designed to span an intron (Brack, *et al.*, *J Clin Invest*, pp.99:2842-2850 (1997)). Amplified products were labeled with digoxigenin-11-dUTP (Boehringer Mannheim, Indianapolis, IN) and then semiquantified in a liquid hybridization assay with biotinylated internal probes (Table 1) using a commercially available PCR ELISA kit (Boehringer Mannheim). In this assay, the labeled PCR products were hybridized with 200 ng/mL probe at 42°C for β -actin, interleukin (IL)-4, IFN- γ , transforming growth factor (TGF)- β 1, and tumor necrosis factor (TNF)- α and at 55°C for IL-1 β and IL-10 for two hours. Hybrids were immobilized on streptavidin-coated microtiter plates and, after washing, were detected with a peroxidase-labeled anti-digoxigenin antibody. Plates were developed by a color reaction using ABTS (2,2'-azino-di[3-ethylbenzthiazoline sulfonate] diammonium salt) substrate and quantitated using a kinetic microplate reader (Molecular Devices, Sunnyvale, CA). The number of cytokine-specific sequences was determined by interpolation with a standard curve and was expressed as the number of cytokine sequences per 2×10^6 β -actin sequences.

Table 1. Nucleotide Sequences of PCR Primers and Biotinylated Probes

		Oligonucleotide sequence
5	β -Actin	5' Primer ATG GCC ACG GCT GCT TCC AGC
		3' Primer CAT GGT GGT GCC GCC AGA CAG
		Probe TAC AGG TCT TTG CGG ATG TC
	IL-1 β	5' Primer GAC ACA TGG GAT AAC GAG GC
		3' Primer GGG ATC TAC ACT CTC CAG CTG
		Probe AGC TTT TTT GCT GTG AGT CCC GGA G
10	IL-4	5' Primer CTT CCC CCT CTG TTC TTC CT
		3' Primer TTC CTG TCG AGC CGT TTC AG
		Probe AGA GCA GAA GAC TCT GTG CAC CGA G
	IL-10	5' Primer CAG TTT TAC CTG GAG GAG
		3' Primer CAA TAA GGT TTC TCA AGG GGC TGG GTC
		Probe CTA CGG CGC TGT CAT CGA TTT CTT
15	IFN- γ	5' Primer ACC TTA AGA AAT ATT TTA ATG C
		3' Primer ACC GAA TAA TTA GTC AGC TT
		Probe ATT TGG CTC TGC ATT ATT TTT CTG T
	TGF- β 1	5' Primer AAG TGG ACA TCA ACG GGT TCA CTA
		3' Primer GCT GCA CTT GCA GGA GCG CAC
		Probe ATC TGC AAA GCT CCC GGC AC
20	TNF- α	5' Primer TAG CCC ATG TTG TAG CAA ACC C
		3' Primer TCG GCA AAG TCG AGA TAG TC
		Probe AAT GGC GTG GAG CTG AGA GAT AAC

25 5. Statistical Analysis

The clinical presentations of the histopathology defined patient subsets were compared by using a Fisher's exact test. *In situ* cytokine production was compared using a nonparametric Kruskal-Wallis test.

6. Results

Microscopic evaluation of rheumatoid synovium revealed fibrin exudation and hyperplasia of synovial lining cells, often assuming a villous configuration, stromal fibrosis, capillary angiogenesis, and diffuse stromal inflammation. Tissue-infiltrating cells included T cells, macrophages, and B cells. Giant cells were variably present. To identify pathological patterns of the synovial inflammation, a series of 21 tissues from patients undergoing joint surgery was analyzed. To exclude features due to long standing and burned out diseases, only patients with clinically active synovitis were enrolled.

Upon analysis by conventional histology, three patterns emerged. Ten tissue samples were categorized as diffuse synovitis (Table 2). These tissues were characterized by a diffuse infiltrate of lymphocytes and macrophages without any additional microanatomical arrangements. The infiltrate tended to be sparse and was accompanied by moderate edema as well as delicate and diffuse fibrosis. A second pattern manifested as demarcated lymphocytic aggregates with sparing of the intervening stroma. In some patients, indistinct germinal center formation with central clearing of the aggregates was apparent. Immunohistochemical analysis showed that follicular structures displayed a central accumulation of B cells surrounded by T cells. Such pseudo-follicular organizations were detected in seven tissues and were classified as follicular synovitis. Four patients displayed necrobiotic granulomas. In this pattern, a fibrinoid necrotic center was lined by a collar of epithelioid histiocytes with or without giant cells. External to this palisade, or garland, of histiocytes was a zone of granulation tissue with angiogenesis and a mixed inflammatory infiltrate composed of lymphocytes, histiocytes (macrophages), and plasma cells. No specimens displayed both follicular synovitis and granulomatous necrobiosis.

Each specimen was evaluated further with respect to fibrinous exudates, capillary neovascularization, and linear subsynovial inflammatory arrays. These features were variable and were not correlated with each other or any of the three main inflammatory patterns.

Table 2. Histomorphological Characteristics of Rheumatoid Synovium

Patient	Diffuse infiltrate	Follicle formation	Granuloma formation
RA 1	+	-	-
RA 2	+	-	-
RA 3	+	-	-
RA 4	+	-	-
RA 12	+	-	-
RA 13	+	-	-
RA 14	+	-	-
RA 15	+	-	-
RA 18	+	-	-
RA 21	+	-	-
RA 7	+	+	-
RA 8	+	+	-
RA 10	+	+	-
RA 16	+	+	-
RA 17	+	+	-
RA 19	+	+	-
RA 20	+	+	-
RA 5	+	-	+
RA 6	+	-	+
RA 9	+	-	+
RA 11	+	-	+

Example 2 - T-cell Derived Cytokines in Distinct Types
of Rheumatoid Synovitis

5 A semiquantitative PCR/liquid hybridization assay was used to detect low concentrations of cytokines in tissue extracts. To directly address the question of whether a correlation exists between the tissue organization of inflammatory cells and cytokine production, the three histomorphologically defined types of synovitis were compared for *in situ* transcription of the IFN- γ and IL-4 genes. To correct for variations in the amount of tissue used for RNA extraction, all cDNAs were adjusted to a concentration equivalent to 2×10^6 copies of the β -actin gene product (Figure 1). IFN- γ mRNA production was a characteristic finding for granulomatous synovitis. Tissues with follicular synovitis contained variable concentrations of IFN- γ mRNA copies (median of 114 copies) whereas the lowest levels were found in extracts derived from material with diffuse synovitis (median of 55 copies). As demonstrated by two-color immunohistochemistry, the major IFN- γ producing cells in all three forms of synovitis were T cells. The high rate of IFN- γ transcription distinguished patients with granuloma formation from the diffuse subtype ($P = 0.007$).

15 A different pattern of tissue distribution was seen for IL-4 mRNA. IFN- γ and IL-4 are both derived from T cells, but they are usually secreted by distinct subsets of T helper cells. T cells with a commitment to the T_H1 pathway release IFN- γ and IL-2. Conversely, IL-4 is the typical product of a T_H2 cell. IFN- γ and IL-4 can be considered as antagonistic effectors with IFN- γ driving inflammatory responses and IL-4 acting as an anti-inflammatory mediator. The highest concentration of IL-4 mRNA was determined in samples with granulomatous synovitis (median of 116 copies). IL-4 transcripts were essentially absent in biopsies from patients with follicular synovitis and were detectable only at low levels in diffuse synovitis.

25 Taken together, the studies on T-cell derived cytokines indicated that diffuse synovitis correlated with low concentrations of IFN- γ and IL-4 mRNA. This pattern might indicate a low degree of activation of predominantly noncommitted T_H0 cells. Follicular synovitis followed the paradigm for germinal centers in that IFN- γ was the dominant T cell product and IL-4 was virtually absent. This type of RA would be best described as a T_H1 mediated response. The constellation of high levels of IFN- γ combined with high transcription of IL-4

30

mRNA found in granulomatous synovitis did not fit the current paradigm. The presence of IL-4 mRNA was surprising and raised the question of how these two antagonistic cytokines can co-exist in the lesions.

Example 3 - Differences in Macrophage Activation Distinguish the Three Variants of Rheumatoid Synovitis

To address the question of whether the different organizational forms of rheumatoid synovitis are associated with differences in the activation of synovial macrophages, the number of IL-1 β and TNF- α transcripts were determined in the tissue extracts. IL-1 β and TNF- α specific sequences were detected in all samples. As shown in Figure 2, concentrations of *in situ* transcribed IL-1 β varied extensively, ranging from 103 to 17,400 copies per 2×10^6 β -actin sequences. Low copy numbers were frequently found in specimens with diffuse synovitis (median of 270 copies), and intermediate levels of IL-1 β transcription were typical for follicular synovitis (median of 2038 copies). This difference was statistically significant ($P = 0.019$). Tissue sections with granuloma formation had the highest rate of IL-1 β mRNA synthesis (median of 10,044 copies), and could be clearly distinguished ($P = 0.005$) from tissues with diffuse infiltrates, which had a low level transcription. The results for TNF- α paralleled the findings for IL-1 β ($r^2 = 0.860$; $P < 0.001$; Figure 2). A low copy number (median of 660 copies) was found in diffusely infiltrated tissue compared with an intermediate level of TNF- α transcription in follicular tissue (median of 1799 copies; $P = 0.04$) and high concentrations in granulomatous synovitis (median of 9568 copies; $P = 0.005$).

Low, intermediate, and high IL-1 β and TNF- α production correlated with IFN- γ transcription in all three disease variants ($r^2 = 0.674$ and $P < 0.001$ for IL-1 β and $r^2 = 0.607$ and $P < 0.001$ for TNF- α compared to IFN- γ). This relationship is compatible with a regulatory role of the T cell product, IFN- γ , in macrophage activation. In this model, rheumatoid granulomas could be regarded as sites of marked T cell and macrophage stimulation. The histiocyte formation characteristic of the granulomas would be consistent with this hypothesis.

Example 4 - Anti-Inflammatory Cytokines in Distinct Variants of Rheumatoid Synovitis

The concentrations of IL-10 and TGF- β 1 mRNA in the tissues were semiquantified. IL-10 transcripts were consistently found (Figure 3). The lowest tissue concentration of IL-10 mRNA correlated with the presence of diffuse synovitis. In these patients, the median copy numbers of cytokine per 2×10^6 β -actin sequences was 76. In contrast, abundance of IL-10 transcripts was a characteristic feature of follicular synovitis. Patients with follicular aggregates synthesized a median of 331 copies of IL-10-specific sequences ($P = 0.04$). Despite the marked stimulation of macrophages in the granulomatous synovitis, IL-10 was present at only low levels (median of 181 copies). Although macrophages are the main producer of IL-10 in the inflamed synovium, macrophage stimulation as indicated by IL-1 β production can thus be differentiated from IL-10 production.

TGF- β 1 is understood as a powerful suppressive cytokine. To explore whether the subdued production of T cell and macrophage products in the diffuse synovitis was correlated with a dominance of TGF- β 1 mRNA, this cytokine was analyzed in all three variants of rheumatoid synovitis. TGF- β 1 sequences could easily be detected in all but one patient. Transcript concentrations ranged from 0 to 19,020 copies (Figure 3). Low levels of TGF- β 1 transcription was a common denominator among patients with diffuse synovitis. They synthesized a median of 3901 copies, a finding that distinguished them from patients with follicular disease (median of 12,636 copies; $P = 0.025$).

Therefore, neither IL-10 nor TGF- β 1 could account for the aborted cytokine response in diffuse synovitis. IL-10, so far recognized as a suppressive cytokine, was transcribed in tissue samples with follicular T-B aggregates and was distinctly low in granulomatous synovitis whereas TGF- β 1 mRNA was found in all three forms of synovitis.

Example 5 - Host Factors Correlate with the Organization and Function of Inflammatory Cells in the Rheumatoid Synovium

Results on tissue cytokine pattern suggested that more than one type of rheumatoid synovitis exists with distinct pathways of inflammation. Phenotypic heterogeneity of RA could be attributed to differences in host factors. To address this issue, the three patient subsets were compared for demographic features and for similarities in clinical presentation.

Also, it could be argued that differences in the histopathological appearance and functional profiles of accumulated cells could be influenced by therapeutic intervention. Treatment given in the last few weeks before harvesting of the tissue could be particularly important in affecting the disease process in the synovial membrane.

The demographic characteristics of the patient cohorts stratified according to the histomorphology of the synovitis are presented in Table 3. Sex, age, and disease duration did not predict which variant of synovitis the patient had developed. All patients had long-standing disease, but no differences were seen that correlated with a specific organizational and functional pattern in the tissue. Patients lacking rheumatoid factor production accumulated among the cohort of individuals found to have diffuse synovitis. The group of patients with diffuse synovitis included in five of ten patients who were seronegative, whereas all patients who had developed follicular synovitis secreted rheumatoid factor ($P = 0.04$). The most intriguing finding was that the generation of the rheumatoid nodules was characteristic for individuals with synovial granuloma formation. All patients with granulomatous synovitis, but none of the patients with follicular synovitis and only one of ten patients with diffuse synovitis, had rheumatoid nodules of the skin. The accumulation of patients with nodular disease in the category of granulomatous synovitis was statistically significant ($P = 0.005$ versus diffuse synovitis; $P = 0.003$ versus follicular synovitis). Pertinent treatment information is summarized in Table 4. All individuals with granulomatous synovitis had been treated with disease-modifying agents (DMARDs) such as hydroxychloroquine, steroids, gold, sulphasalazine, azathioprine, and methotrexate. None of these patients had been managed with nonsteroidal anti-inflammatory drugs (NSAIDs) alone. The group of patients with diffuse synovitis appeared to have been treated less aggressively. One-third of these patients was managed with NSAIDs only. All patients with granuloma formation in the synovia, but only 30 and 29%, respectively, of patients in the diffuse and follicular categories, were on methotrexate. Despite the low number of individuals analyzed, these differences showed a trend toward significance ($P = 0.07$ and $P = 0.06$, respectively). This analysis suggested that host factors and/or treatment may contribute to the microanatomy in the joint or *vice versa* (i.e., that a correlation exists between systemic manifestations and certain types of rheumatoid synovitis).

Table 3. Characteristics of Patient Populations Defined by Histopathological Patterns

	Sex (♀/♂)	Median age in years (range)	Median disease duration in years (range)	Rheumatoid factor positive*	Rheumatoid nodules**
Diffuse synovitis	5/5	57 (52-73)	16.5 (1-51)	50%	10%
Follicular synovitis	6/1	57 (34-67)	10 (4-50)	100%	0%
Granulomatous synovitis	3/1	61.5 (24-68)	8 (5-32)	75%	100%

* $P = 0.004$ for diffuse *versus* follicular synovitis.

** $P = 0.005$ for diffuse *versus* granulomatous and $P = 0.003$ for follicular *versus* granulomatous synovitis.

All other comparisons were nonsignificant.

Table 4. Past and Current Treatment in the Different Patient Categories

	Diffuse synovitis (A)	Follicular synovitis (B)	Granulomatous synovitis (C)	p		
				A vs B	A vs C	B vs C
NSAID only (last 3 months before surgery)	4/10	1/7	0/4	NS	NS	NS
DMARD (last 3 months before surgery)	6/10	6/7	4/4	NS	NS	NS
Methotrexate (last 3 months before surgery)	3/10	2/7	4/4	NS	0.07	0.06
DMARD (total)	7/10	6/7	4/4	NS	NS	NS

NSAID, nonsteroidal anti-inflammatory drugs; DMARD, disease-modifying anti-rheumatic drugs; NS, not significant.

Example 6 - Tissue Cytokine Patterns Distinguish Variants of RA

RA is a chronic inflammatory disease with profound phenotypic variability (Harris ED (ed): Rheumatoid Arthritis. Philadelphia, WB Saunders Co., pp.3-212 (1997)). The pattern

of involvement, the course and destructive potential of the disease, and the frequency of extra-articular manifestations vary significantly. Reasons for the phenotypic heterogeneity are not completely understood but may include variable combinations of disease risk genes (Ollier WE, MacGregor A: *Br Med Bull*, pp.51:267-285 (1995); and Weyand CM, Goronzy JJ: *Med Clin North Am*, pp.81:29-55 (1997)). The results presented herein indicate that the heterogeneity of the disease process includes the synovial lesion and that patients display considerable differences in the organization and the functional commitment of the inflammatory infiltrates. The microanatomy of the inflamed synovium showed a correlation with profiles of tissue cytokines, supporting the model that different mechanisms are functional in regulating rheumatoid synovitis. Tissue destruction and possible other aspects of the disease process are related to cell-cell interactions in the infiltrates. Thus, understanding the rules underlying the emergence of a defined microanatomical structure in the synovial membrane provides information about the fundamentally important pathological events leading up to RA.

The results from the study presented herein indicate that the topographical arrangement of the mononuclear infiltrate can be used to define three variants of rheumatoid synovitis. These three patterns correlated with the combination and the amount of cytokines produced in the tissue. In general, the level of transcription of T-cell derived cytokines is low in synovial tissue, and the study thus employed a semiquantitative PCR approach. This technique can have limitations that need to be considered when interpreting the data. As it is a semiquantitative method, differences have to be large to be distinguished. Also, the level of cytokine mRNA may not necessarily reflect the amount of functionally active protein, particularly in the case of TGF- β 1 and IL-1 β . Cytokine production was confirmed by immunohistochemistry, which, however, is not a suitable technique to determine quantitative differences. The finding that 21 samples were sufficient to accomplish a dissection of the three variants emphasizes that the differences in cytokine production were pronounced.

The first variant of RA emerging from the study is a disease phenotype characterized by diffuse infiltrates in the synovia, a lower probability of rheumatoid factor production, and a clinically milder disease that is responsive to nonaggressive treatment. The low transcription of proinflammatory mediators of a T_H0 pattern suggested that the T cell response

was not highly differentiated. Consistent with the interpretation is the absence of a microatomical organization of the mononuclear infiltrate. The reason for this subdued activation of inflammatory cells is unclear. Production of the anti-inflammatory cytokines IL-10 and TGF- β were generally low, and no evidence was found for an activate suppressive mechanism.

The second variant of RA distinguished in this study represents the histomorphological pattern that is typically associated with RA, the formation of follicular structures composed of T and B cells. Recent molecular characterization of B cells isolated from RA tissues with follicular centers have confirmed that these T-B cell aggregates functionally resemble germinal centers (Schroeder, *et al.*, *Proc Natl Acad Sci USA*, pp.93:221-225 (1996); and Randen, *et al.*, *Scan J Immunol*, pp.41:481-486 (1995)). Germinal centers are the site of B cell differentiation, somatic mutation, and affinity maturation, all of which are T-cell-dependent processes.

The tissue cytokine profile that emerged for this category included intermediate levels of IFN- γ with essentially absent IL-4 transcription. This pattern would suggest a T_H1 deviation of the immune response. The predominance of a T_H1 pattern in patients with follicular synovitis is in line with current knowledge in cytokine production in germinal centers (Kelsoe G: *Semin Immunol*, pp.8:179-184)). Although IL-4 deficient mice generated by gene targeting are not able to form lymphoid follicles (Kuhn, *et al.*, *Science*, pp.254:707-710 (1991)), T cells accumulating in germinal centers typically do not produce IL-4 but IFN- γ . Also the production of IL-10 found in the tissues with follicular aggregates may relate to germinal center formation (Levy Y, Brouet JC, *J Clin Invest*, pp.93:424-428)). IL-10 can be produced by a variety of cells, including T_H1 and T_H2 type T-cells, macrophages, and B cells. It acts by inhibiting the production of numerous pro-inflammatory monokines and be attenuating T_H1 mediated immune responses. In contrast to this immunosuppressive effect, IL-10 as marked stimulatory effects on B cells and supports B cell proliferation and differentiation. In particular, the production of rheumatoid factor appears to be IL-10 dependent (Perez, *et al.*, *Arthritis Rheum*, pp.38:1771-1776 (1995)). The data herein would suggest that the production of IL-10 in follicular synovitis relates to providing a microenvironment for B cell proliferation and not for suppressing macrophage activation.

004347 0002260
a T_H1 response, and the use of IL-4 and IL-10 have been proposed as therapeutic interventions (Heinzel, *et al.*, *J Exp Med*, pp.169:59-72 (1989); Yamamura, *et al.*, *Science*, pp.254:277-279 (1991); Simon, *et al.*, *Proc Natl Acad Sci USA*, pp.91:8562-8566, (1994); Van Roon, *et al.*, *Arthritis Rheum*, pp.39:829-835 (1996); Isomaki, *et al.*, *Arthritis Rheum*, pp.39:386-395 (1996); Walmsley, *et al.*, *Arthritis Rheum*, pp.39:495-503 (1996)). The present finding that RA encompasses different cytokine patterns indicates that the choices of the host in terms of cytokine recruitment do not determine whether or not the individual develops the disease but can influence disease severity and organ involvement.

Alternatively, the three disease phenotypes may reflect distinct pathomechanisms including different disease initiators. As the initial events in RA are not understood, both models remain feasible. However, searches into the instigators of RA could benefit from the realization that there exist several variants of the disease. It might be misleading to search for common denominators in a cohort of RA patients if multiple distinct disease variants are represented. Focusing on a single entity of RA may enhance the identification of the shared pathomechanisms, genetic risk factors, and antigens driving rheumatoid synovitis. Equally important, phenotypic variants of RA should be considered in the design of treatment trials and in the application of therapeutic agents in individual patients.

In summary, RA is a chronic inflammatory disease with primary manifestations in the synovial membrane. Tissue infiltrates are composed of T cells, B cells, and macrophages, but histopathological appearances vary widely and rarely pathognomonic. Mechanisms underlying the phenotypic heterogeneity of rheumatoid arthritis are not known. To explore whether a correlation exists between the microscopic patterns of rheumatoid synovitis and *in situ* production of cytokines, tissue samples from 21 consecutive patients with clinically advanced active RA were examined. Based upon the organization of the lymphocyte infiltrate, the synovial biopsies were categorized into three distinct subsets. Ten samples were characterized by diffuse lymphoid infiltrates without further microarrangements. In seven samples, lymphoid follicles with germinal center formation were detected, and in four specimens, granuloma formation was identified. In all specimens, cytokine transcription of interferon (IFN)- γ , interleukin (IL)-4, IL-1 β , tumor necrosis factor (TNF)- α , IL-10, and transforming growth factor- β 1 was semiquantified with polymerase chain reaction and liquid

participate in this type of granuloma formation is not entirely clear. For example, it is not understood what the macrophage-activating agent in this type of lesion is. Nevertheless, T_H1 as well as T_H2 associated cytokines can apparently participate in granuloma formation.

The granuloma formation in the rheumatoid synovium was different from these experimental models in that IFN- γ and IL-4 were co-produced. Previous results have emphasized their antagonistic effects (Seder RA, Paul WE: *Annu Rev Immunol*, pp.12:635-673 (1994); Abbas. *et al.*, *Nature*, pp.383:787-793 (1996); Seder, *et al.*, *J Exp Med*, pp.176:1091-1098 (1992); Seder, *et al.*, *Proc Natl Acad Sci USA*, pp.90:10199-10192 (1993); and Szabo, *et al.*, *J Exp Med*, pp.185:817-824 (1997)). Neutralization of IFN- γ production enhances granuloma formation in the schistosoma model although it abrogates the inflammation in the microbacterial model. However, it is possible that these findings are restricted to the early stages of granuloma formation and many not apply to chronic inflammation, as is the case in the rheumatoid synovium. Indeed, a temporal participation of T_H1 as well T_H2 cells has been described for the mycobacterial infection as well as for the schistosoma-egg-induced granuloma (Boros DL, *Immunobiology*, pp.191:441-450 (1994); Chensue, *et al.*, *J Immunol*, pp.151:1391-1400 (1993); and Orme, *et al.*, *J Immunol*, pp.151:518-525 (1993)).

The data presented herein indicate the contribution of different cytokines in controlling the microanatomy of the synovial inflammation and also emphasize the necessity to re-evaluate the role of the immune pathways in the pathogenesis of RA. Immune deviation is now widely accepted as a concept in explaining the pathogenesis of autoimmune diseases. This model implies that chronic inflammation is a consequence of the aberrant commitment to an immune pathway in response to a given antigen (Finkelman FD, *J Exp Med*, pp.182:279-282 (1995). Studies in diseases such as infection with leishmania major and mycobacterium leprae have fueled the hypothesis that immune reactions to a given antigen can take very different paths, and accordingly, a certain type of disease may develop (Heinzel, *et al.*, *J Exp Med*, pp.169:59-72 (1989); Yamamura, *et al.*, *Science*, pp.254:277-279 (1991)). Tuberculoid leprosy and lepratomatous leprosy have been distinguished as two disease phenotypes, the distinction being mainly attributed to the involvement of different cytokine networks. Following this model, RA has been considered a consequence of an immune deviation toward

phase hybridization. Each of the morphologically defined variants of synovitis displayed a unique cytokine profile. Low-level transcription of IFN- γ , IL-4, IL-1 β , and TNF- α was typical of diffuse synovitis. In follicular synovitis, IFN- γ was the dominant cytokine, IL-4 was virtually undetectable, and IL-10 was abundant. Granulomatous synovitis demonstrated high transcription of IFN- γ , IL-4, IL-1 β , and TNF- α and could be clearly distinguished from the other phenotypes. To investigate whether differences in the synovial lesions were related to host factors, patients were compared for clinical parameters. Diffuse synovitis was seen in most of the patients with seronegative RA, the mildest form of the disease. In contrast, extra-articular spreading of RA with nodule formation was typically associated with granulomatous synovitis.

Example 7 - HLA-DRB1 alleles associated with RA

1. The shared epitope hypothesis

The association of RA to MHC class II genes was recognized in the 1970s and provided the framework for an important disease model (Rittner, *et al.*, *Mol Med* 3:452 (1997)). Sharing of MHC class II alleles in RA patients was taken as evidence that the major biological function of these molecules, the presentation of antigenic peptides to T cells, was critically involved in disease pathogenesis (Todd, *et al.*, *Science* 240:1003 (1988)). A widely accepted paradigm holds that RA represents the sequel of pathologic T cell responses initiated and maintained by antigens presented in the context of disease-associated HLA molecules. The initial observation that the frequency of HLA-DR4 was increased in RA patients was confirmed in multiple studies and in multiple ethnic groups. The association with HLA-DR4 was eventually attributed to some of the allelic variants of the HLA-DR4 family, including HLA-DRB1 *0401, *0404, *0405, and *0408 (Table 5). Sequence similarity between these disease-associated alleles and the sequence differences to the nonassociated allele HLA-DRB1 *0402 gave rise to the "shared epitope" hypothesis (Gregersen, *et al.*, *Arthritis Rheum* 30:1205 (1987); Nepom, *et al.*, *Annu Rev Immunol* 9:493 (1991); and Winchester R., *Adv Immunol* 56:389 (1994)). This shared sequence stretch was also identified in HLA-DRB1 *01 alleles enriched in Jewish patients with RA and in the HLA-DRB1 *1402 allele in Yakima Indians (Willkens, *et al.*, *Arthritis Rheum* 34:43 (1991); and Willkens, *et al.*,

associated alleles that influence the binding properties of several HLA-DR pockets, but these are not considered to be critical for the disease association. Disease-associated HLA-DR4 variants have either a Val or a Gly at position 86. This dimorphism has a major impact on T cell recognition, probably by altering the specificity of peptide binding pocket 1 (Fu, *et al.*, *J Exp Med* 181:915 (1991)). The collagen peptide crystallized with HLA-DRB1 *0401 has a Met in position 1 and is therefore unlikely to bind to HLA-DRB1 *0404. Also, the size of pocket 4 is smaller in DRB1 *0404/8 (Arg at position 71) than in DRB1 *0401 (Lys at position 71). Sequence polymorphisms in HLA-DR1 alleles add additional complexity. Amino acids at position 13, which are different in HLA-DR1 and HLA-DR4, contribute to binding pocket 4. Pocket 6 is deeper in DR4 than in DR1 because of the smaller side chain of Val in DR4 at position 11 compared with the Leu in DR1. In contrast, a Tyr versus Ser exchange at position 30 reduces the size of pocket 7 in DR4 in comparison with DR1. Additional different binding properties can be predicted for other disease-associated HLA-DRB1 alleles, such as DRB1 *1402 and *1001.

Alternatively, models have been proposed to explain the sharing of a sequence polymorphism of the HLA-DRB1 molecule in RA. Carson *et al.* have favored the hypothesis that the peptide encoded by the shared epitope is recognized as a self-antigen and may therefore influence thymic selection (Roudier, *et al.*, *Proc Natl Acad Sci U S A* 86:5104 (1989)). Support for this model has come from subsequent studies demonstrating that HLA-DR-derived peptides can be presented in restriction to HLA-DQ (Salvat, *et al.*, *J Immunol* 153:5321 (1994)). Albani and Carson have shown that the recognition of microbial peptides expressing sequence homologies to the shared epitope is altered in RA patients (Albani, *et al.*, *Immunol Today* 17:466 (1996)). Based on these data, this group has developed a molecular mimicry model of RA in which positive selection in the thymus favors T cells that cross-react between microbial and self-antigens (Albani, *et al.*, *Nat Med* 1:448 (1995)). An alternative hypothesis has been proposed by David *et al.* who also predict a role of HLA-DR-derived peptides in thymic selection (Taneja, *et al.*, *J Clin Invest* 101:921 (1998)). They, however, have postulated that peptides derived from non-associated HLA-DR molecules, and not from disease-associated molecules, are functionally important and that these peptides somehow select regulatory T cells that protect against collagen-induced arthritis.

Finally, it is possible that the shared epitope functions in RA by directly contacting the TCR molecule. This model has been originally proposed based on response patterns and TCR BV gene segment usage of alloreactive T cell clones and the mapping of antibody binding sites on the HLA-DR β 1 chain by site-directed mutagenesis (Goronzy, *et al.*, *Rheum Dis Clin North Am* 21:655 (1995); Hiraiwa, *et al.*, *Proc Natl Acad Sci U S A* 87:8051 (1990); and Weyand, *et al.*, *Arthritis Rheum* 37:514 (1994)). More recently, the crystal structure of HLA-DR peptide complexes has confirmed that amino acids at position 70 of the HLA-DR β 1 chain project out of the binding site and may directly interact with the TCR (Dessen, *et al.*, *Immunity* 7:473 (1997); and Stern, *et al.*, *Nature* 368:215 (1994)). Penzotti *et al.* (*Arthritis Rheum* 40:1316 (1997)) have modeled a TCR from an HLA-DR4-specific clone and the DRB1*0404 molecule and have predicted that TCR residues CDR1 β Asp30, CDR2 β Asn51, and CDR3 β Gln97 are positioned to participate in hydrostatic interactions with the DRB1 residues Q and R at positions 70 and 71, respectively, of the shared epitope. These contact residues are likely important in repertoire selection. Indeed, the RA-associated HLA-DR polymorphism has been found to induce global repertoire changes that are reflected at the level of TCR BV-BJ gene segment frequencies (Walser-Kuntz, *et al.*, *Immunity* 2:597 (1995)). In particular, TCRs were affected that had a consensus Gly-Pro-Gly sequence and therefore were predicted to have a more rigid structure.

Table 5. HLA-DR association of RA

Serotype	Cellular type	Genotype	Disease association
HLA-DR4	HLA-Dw4	HLA-DRB1 *0401	+
	HLA-Dw10	HLA-DRB1 *0402	-
	HLA-Dw13	HLA-DRB1 * 0403	-
		HLA-DRB1 *0407	-
	HLA-Dw14	HLA-DRB1 *0404	+
		HLA-DRB1 *0408	+
	HLA-Dw15	HLA-DRB1 *0405	+
HLA-DR1	HLA-Dw1	HLA-DRB1 *0101	+

HLA-DRw6	HLA-Dw9	HLA-DRB1 *1401	-
	HLA-Dw16	HLA-DRB1 *1402	+
	HLA-Dw18	HLA-DRB1 *1301	-
	HLA-Dw19	HLA-DRB1 *1302	-

5

2. The alternative view - A "multiple HLA polymorphisms - multiple disease variants" model

The HLA-DR association of RA has been studied in many ethnic groups, and all disease-associated alleles defined so far, express the "shared epitope." While this observation suggests a common denominator and a single pathomechanism underlying the HLA association of RA, several aspects of these studies require a rethinking of such a simplified model. First, it should be noted that the strength of the association is not uniform but varies widely depending on the ethnic background of the patients. Second, not all shared-epitope alleles are equal, for example, HLA-DRB1*0401 may have an exceptional role.

Indications that this genetic heterogeneity may be useful in dissecting the clinical heterogeneity of RA came from association studies in Caucasian patients stratified for different disease manifestations (Weyand, *et al.*, *Ann Intern Med* 117:801 (1992); and Weyand, *et al.*, *J Clin Invest* 89:2033 (1992)). Earlier studies had indicated that HLA-DR4 may be associated with more erosive disease (Olsen, *et al.*, *Am J Med* 84:257 (1988)). Subsequent studies showed that patients with rheumatoid factor (RF)-negative (seronegative) RA were characterized by an enrichment of both the HLA-DRB1*04 and *01 alleles (Weyand, *et al.*, *J Clin Invest* 95:2120 (1995)). In contrast, 90% of patients with erosive seropositive disease had at least one copy of an HLA-DRB1*04 allele. Sequence analysis of HLA-DRB1*04 genes in the seronegative cohort identified the DRB1*0404 and B1*0408 variants as the dominant alleles. This contrasted to the overrepresentation of B1 *0401 in the seropositive patients. Clinical dissection of the seronegative patients demonstrated that erosive disease occurred more frequently in the HLA-DRB1*04-positive individuals. A similar correlation between the inheritance of HLA-DRB1 alleles and the severity of joint destruction emerged for patients with seropositive RA. While all of these patients had erosive disease, patients carrying two disease-associated alleles, either HLA-DRB1*01/04 or HLA-

DRB1*04/04, had a higher frequency of joint arthroplasty (Weyand, *et al.*, *Ann Intern Med* 117:801 (1992)). Based on these observation, it is now proposed that there is a hierarchy of HLA-DRB1 polymorphisms determining the severity of joint inflammation. This model would imply that the disease-associated HLA-DRB1 polymorphism is not only functioning in disease initiation but also in disease progression. As described herein, HLA-DRB1 typing can be used as a prognostic marker to predict disease severity.

An alternative approach would suggest that the clinical phenotypes of RA are not simply steps on a severity scale. Rather, they could be viewed as distinct entities of the disease with ultimately unique pathogeneses. Disease-associated HLA-DRB1 alleles, in spite of sharing a similar sequence stretch, may differ in their contribution to the disease. This model also includes that more than one inflammatory pathway can lead to synovitis and that the clinical category of polyarthritis is a "mixed bag" with multiple forms of inflammation. Assigning different genotypes to different types of RA also provides an explanation for the finding that the strength of association between HLA-DR4 and disease is not maintained in all populations. The majority of African-American patients lack HLA-DR4 (McDaniel, *et al.*, *Ann Intern Med* 123:181 (1995)). While this finding could raise the question whether HLA class II molecules are a prerequisite in the pathological events typical for the disease, it could also be interpreted as demonstrating fundamental differences in the disease processes leading to symmetrical polyarthritis in Caucasians and African-Americans. Similar considerations could be applied to population-based studies describing a lack of association with HLA-DR4 in patients diagnosed with symmetric polyarthritis.

Some clinical observations reinforce that the different disease patterns encountered in RA may actually represent different entities. Impressive discrepancies in the disease course of synovial inflammation and extra-articular RA are compatible with independent disease mechanisms in these two different dimensions of RA. Major organ manifestations frequently occur in RA patients at times when the synovial disease is quiescent. In selected patients, rheumatoid nodules involving the skin or other organs can be induced by methotrexate although the treatment successfully controls synovial inflammation (Merrill, *et al.*, *Arthritis Rheum* 40:1308 (1997)). Rheumatoid nodules impress histomorphologically as a granulomatous reaction, distinct from the inflammatory lesions of rheumatoid synovitis. It

has, therefore, been suggested that extra-articular disease is a different dimension of RA with different pathomechanisms and not just a more severe form of the disease. In Caucasians, major organ involvement is essentially restricted to patients who bear HLA-DR4 and it is generally not seen in patients who express only one of the other disease-associated alleles, again emphasizing the unique role of the HLA-DR4 haplotype. Even more striking is a gene-dose effect in patients with extra-articular manifestations. Patients with erosive RF-positive joint disease are characterized by the genotype HLA-DRB1*04/x (x represents an RA non-associated allele). In contrast, nodule formation is a hallmark of patients with the genotypes DRB1*01/04 or DRB1*0401/0404 (Weyand, *et al.*, *J Clin Invest* 89:2033 (1992)).

Homozygosity for DRB1*0401 is a strong predictor for rheumatoid vasculitis and Felty's syndrome (Lanchbury, *et al.*, *Hum Immunol* 32:56 (1991); and Weyand, *et al.*, *J Clin Invest* 89:2033 (1992)) both of which are severe complications of RA associated with significant morbidity and mortality. The importance of a gene-dose effect in extra-articular disease does not necessarily imply an additive effect of RA-associated HLA alleles on a severity scale. A synergistic interaction between the two haplotypes is a more likely scenario. In this model, the functional role of HLA-DRB1 alleles in synovial and extra articular disease is regarded as distinct, e.g., peptide selection in synovitis and repertoire formation in extra-articular disease.

In summary, the diagnostic category of RA is an umbrella term that encompasses multiple subtypes, each of which is associated with different HLA-DR polymorphisms. As the heterogeneity of the disease should have a major impact on disease course and therapeutic responsiveness, HLA-DR molecules could be used to dissect the crude category of RA into pathogenically homogeneous subsets.

3. Patient population and HLA-DRB1 genotyping

The HLA-DRB1 genotype of patients classified as having a diffuse, follicular, or granulomatous phenotype was determined using PCR according to the methods and materials provided by One Lambda (Los Angeles, CA) or Biotest Diagnostics (Denville, NJ).

Example 8 - Disease phenotypes in RA and genetic control of T cell function

Additional evidence for distinct pathomechanisms in different forms of RA has come from studies of abnormal T cell functions in RA. Several lines of evidence have implicated T cells as important players in RA pathogenesis. The MHC class II association has emphasized the critical role of TCR-MHC interactions in the pathogenesis of the disease (Winchester R., *Adv Immunol* 56:389 (1994)). Equally important is the histomorphology of rheumatoid synovitis, which points to T cells as relevant component of pathogenic events. Inflammatory lesions in the synovial membrane are composed of T cells, macrophages, B cells, and synoviocytes. Notably, these inflammatory cells acquire distinct topographic arrangements of which follicular T cell-B cell aggregates are the most interesting (Kurosaka, *et al.*, *J Exp Med* 158:1191 (1983)). Follicular structures in rheumatoid synovitis display morphologic, phenotypic, and functional characteristics resembling germinal centers (Schroder, *et al.*, *Proc Natl Acad Sci U S A* 93:221 (1996)). The formation of germinal centers, usually restricted to lymphoid organs, is a T cell dependent process and provides a unique microenvironment for the generation of high affinity B cell responses. Most T cell studies in RA have therefore focused on identifying and characterizing the TCRs that might recognize antigen in the synovial tissue. Repertoire analysis of synovial T cells have demonstrated that the infiltrate is heterogeneous, that some T cells are clonally expanded, and that different T cells are dominant in different patients. These studies have emphasized the heterogeneity of the disease process without defining a common denominator. Relatively few studies have attempted to identify variables of T cell function that might be genetically determined and reflect disease-risk genes. Allelic polymorphisms of TCR genes have received the greatest attention, but these studies have remained inconclusive (Cornelis, *et al.*, *Arthritis Rheum* 40:1387 (1997); and Hall, *et al.*, *Arthritis Rheum* 40:1798 (1997)). More recent data, however, indicate that abnormalities in the T cell repertoire and some unexpected T cell functions in RA have a genetic basis and correlate with the clinical expression of RA.

Several groups have made the observation that RA patients carry clonally expanded T cell populations not only in the synovial compartment, as one would expect, but also in the peripheral T cell repertoire (DerSimonian H, *et al.*, *J Exp Med* 177:1623 (1993); Fitzgerald JE, *et al.*, *J Immunol* 154:3538 (1995); Gonzalez-Quintial R, *et al.*, *J Clin Invest* 97:1335

detected. Rather, the expression of CD28-deficient CD4 T cells is a hallmark of early and untreated disease. Most importantly, indirect evidence suggests genetic control of CD4⁺/CD28^{null} T cells. A study evaluating the presence of CD4⁺/CD28^{null} T cells in monozygotic and dizygotic twins and in spouse pairs revealed a high concordance in the monozygotic twin pairs but a lack of correlation in the spouse pairs. It is proposed herein that the generation of CD4⁺/CD28^{null} T cells is at least partially influenced by inherited factors.

The molecular basis of CD28 deficiency has been addressed as the loss of this important costimulatory molecule and may provide clues as to how these cells emerge. The expression of CD28 on the cell surface has been correlated with the presence of two DNA binding proteins that bind to sequence motifs within the minimal promoter of the CD28 gene (Vallejo AN, *et al.*, *J Biol Chem* 273:8119-29 (1998)). These regulatory proteins are absent in CD4⁺/CD28^{null} T cells, providing a tool to approach the mechanisms controlling CD28 expression. So far, besides RA, one more factor has been associated with the loss of these gene specific regulatory proteins, namely advanced age of the donor.

As outlined, the presence of CD4 T cells defective for CD28 expression correlated with the presence of extra-articular disease. Available data indicate that CD4⁺/CD28^{null} T cells differ from traditional CD4⁺ helper T cells in several aspects. CD4⁺/CD28^{null} T cells are dependent on costimulatory signals provided by accessory cells to be fully activated (Park W, *et al.*, *Eur J Immunol* 27:1082 (1997)). However, blocking studies with CTLA-4-Ig fusion protein have demonstrated that costimulation does not involve a CD28/CTLA-4-CD80/CD86 interaction. Also, costimulation enhances the proliferative response and upregulates IL-2R expression but has no effect on the cytokine production. Compared to CD4⁺/CD28⁺ T cells, CD28-deficient CD4⁺ T cell clones produce higher amounts of IFN- γ but similar amounts of IL-2 and IL-4. IFN- γ production is not enhanced by accessory cells, suggesting that the secretion of IFN- γ by CD4⁺/CD28^{null} T cells is independent of costimulatory mechanisms. As a second important difference to traditional helper cells, CD4⁺/CD28^{null} T cells lack the expression of CD40 ligand (CD40L) and fail to provide helper cell signals for B cell differentiation and immunoglobulin production (Weyand CM, *et al.*, *Mech Age Develop* In press (1998)). In contrast, they have been found to transcribe and produce perforin, a molecule tightly linked with cytotoxic capability (Namekawa T, *et al.*, *Arthritis Rheum.*

(1996); Goronzy JJ, *et al.*, *J Clin Invest* 94:2068 (1994); Hingorani R, *et al.*, *J Immunol* 156:852 (1996); and Lim A. *et al.*, *Hum Immunol* 48:77 (1996)). Genetic susceptibility for the formation of T cell oligoclonality was suggested by the finding that CD4 T cell clones were present in affected as well as unaffected members of multi-case families with RA (Waase I, *et al.*, *Arthritis Rheum* 39:904 (1996)). The isolation of clonally expanded CD4 T cells from patients with RA has allowed for the phenotypic and functional characterization of these T cells (Schmidt D, *et al.*, *J Clin Invest* 97:2027 (1996)). Expanded CD4 clonotypes were found to be deficient for the expression of the CD28 molecule. Because CD28 has been identified as a key player in T cell costimulation, this finding raised the question whether CD28-deficient CD4 T cells are functionally competent and how, if so, they contribute to RA. CD28-deficient CD4 T cells, as opposed to CD8⁺/CD28^{null} T cells, are infrequent in normal individuals. Most normal donors have <1% CD4⁺/CD28^{null} T cells (Martens PB, *et al.*, *Arthritis Rheum* 40:1106 (1997)). In contrast, increased frequencies of CD28-deficient CD4 T cells are detected in patients with RA (Figure 4). Association studies of CD4⁺/CD28^{null} T cells in patient cohorts stratified for different disease manifestations showed that the size of the CD4⁺/CD28^{null} T cell compartment is closely associated with extra-articular RA (Figure 5). Patients with RA limited to the joint expressed a median of 1.1% CD4⁺/CD28^{null} T cells. In patients with rheumatoid nodules, 7% of the CD4 T cells were CD28-deficient. CD4 T cells lacking CD28 expression were encountered in almost all patients with rheumatoid organ disease. In patients with rheumatoid vasculitis, CD28-deficient T cells accounted for up to 50% of the CD4⁺ subset. A role of CD4⁺/CD28^{null} T cells in extra-articular aspects of RA is also supported by their distribution pattern. These unusual CD4 T cells are circulating in the blood and are generally encountered in the synovial tissue, but they are not selectively enriched in the synovial compartment (Rittner HL, *et al.*, *Mol Med* 3:452 (1997)).

It could be argued that the emergence of CD4⁺/CD28^{null} T cells in RA patients is a consequence of chronic persistent inflammation. Several lines of evidence do not support this interpretation. In the normal population, "carriers" for CD4⁺/CD28^{null} T cells can be found, although they are infrequent. In these normal individuals, as well as in RA patients, the proportion of CD4⁺/CD28^{null} T cells remains constant over several years. No influence of disease duration or therapy on the expansion of the CD4⁺/CD28^{null} compartment has been

- Production of high concentrations of IFN- γ
- Autoreactivity in response to adherent cells
- Impaired apoptosis with overexpression of *bcl-2*
- Transcriptional block of the CD28 gene due to a deficiency of two gene-specific transcription factors

1. Cell surface staining to determine CD4⁺/CD28^{null} cell frequency

PBMC were isolated from the patient and stained (20 minutes at 4°C) with a combination of two of the following monoclonal antibodies, anti-CD3 (fluorescein isothiocyanate [FITC]-conjugated), anti-CD4 (peridinin chlorophyll protein [PerCP]-conjugated), anti-CD8 PerCP-conjugated (all from Becton Dickinson, San Jose, CA), and anti-CD28 FITC-conjugated (Pharmingen, San Diego, CA), and analyzed on a FACS Calibur flow cytometer (Becton Dickinson) to determine the frequencies of CD3⁺CD4⁺, CD3⁺CD8⁺, and CD4⁺CD28^{null} T cells. Analysis was performed using WinMIDI software (Joseph Trotter, Scripps Research Institute, La Jolla, CA).

2. Flow cytometric analysis

Twenty thousand events were analyzed using WinMDI software. A tight light scatter region was drawn to include only viable lymphocytes. CD3⁺CD4⁺, CD3⁺CD8⁺, CD4⁺CD28⁺, CD4⁺CD28^{null} cells were analyzed for their expression of the cytokines of interest. In all experiments, less than 1% of the cells were positive for the isotype controls, demonstrating a very high staining specificity.

Example 9 - Variants of RA defined by distinct patterns of synovial inflammation

The primary manifestations of RA are found in the synovial membrane of diarthroidal joints. Histopathologic changes include dense infiltrates of mononuclear cells, neoangiogenesis, and hypertrophy and hyperplasia of synoviocytes. Although it is reasonable to assume that all of the different cell types accumulated in the synovial lesions have a role in the pathologic events, the exact contributions of monocytes, T cells, B cells, and resident

synovial cells have not been delineated. Keeping in mind that more than one form of RA may exist creates the challenging question whether the contribution of these different cells to the inflammation may be variable. Heterogeneity of rheumatoid synovitis is certainly encountered as far as the histomorphologic appearance of the disease lesions is concerned.

5 Pathologic findings in synovial tissues have so far not been helpful in guiding clinical management. While most other fields in medicine have used more sophisticated approaches in tissue diagnostics, this has not been the case for rheumatology. RA has remained primarily a clinical diagnosis (Arnett FC, *et al.*, *Arthritis Rheum* 31:315 (1988)). It is possible that the presumed lack of information gained from histomorphologic inspection of synovial lesions
10 relates to the misleading expectations that RA is a narrowly defined category that should produce a unique pathomorphology.

Evidence for heterogeneity of rheumatoid synovitis comes from the data presented herein correlating the microanatomy of the inflamed synovial membrane with tissue cytokine patterns. T cells, B cells, and macrophages can acquire a highly specialized topography in the
15 synovium. These cells usually accumulate in the sublining stroma and are either arranged as diffuse infiltrates or are organized into structures reminiscent of secondary lymphoid tissues. One of the remarkable structures found in rheumatoid synovitis is T cell-B cell aggregates, which form germinal centers (GCs). Synovial GCs include all of the necessary cellular components and the defined topographical arrangement required for GC reactions to occur.
20 Specifically, networks of CD23⁺ follicular dendritic cells are localized in the centers of T cell-B cell follicles. These networks are populated by IgD⁺ B cells and surrounded by CD4⁺ T cells, some of which express CD40L (Wagner UG, *et al.*, *J Immunol.* 161:6390-7 (1998)). The formation of GCs in the synovial membrane, a primarily non-lymphoid tissue, is a highly significant event that emphasizes the role of the immune system in this disease and stresses
25 the critical contribution of antigen and antigen recognition in pathology. It is, however, important to note that only a subset of patients forms such synovial GCs. Histopathologic patterns of rheumatoid synovitis were examined in a series of biopsies collected from patients with active arthritis (Klimiuk PA, *et al.*, *Am J Pathol* 151:1311 (1997)). In this case series, three distinct patterns emerged. One-third of all samples contained GC-like follicles. In a
30 slightly higher proportion of samples, T cells, B cells, and CD68⁺ macrophages lacked a

specialized microanatomical formation, but formed a diffuse infiltrate. A third pattern of synovitis was encountered in the minority of samples. This granulomatous pattern was characterized by necrotic centers lined by a collar of epithelioid histiocytes, with or without giant cells. Typically, marked angiogenesis was found external to the palisade of histiocytes in areas with mixed infiltrates composed of lymphocytes, plasma cells, and macrophages. It is important to note that in this study, follicular synovitis and granulomatous necrobiosis did not co-occur. Rather, patients appeared to either organize the synovial lesions into follicular or granulomatous patterns.

To examine the relevance of microanatomical arrangements for the inflammatory process, tissue cytokine profiles were determined. IFN- γ and IL-4 were chosen as the marker cytokines for T_H1 and T_H2 helper cells. IFN- γ and IL-4 were present in low abundance in tissues with diffuse synovitis. In follicular synovitis IFN- γ was transcribed at intermediate levels but IL-4 was distinctly absent. Granulomatous synovitis was associated with the highest tissue concentrations for both IFN- γ and IL-4. The production of the macrophage products, IL-1 β and TNF- α , correlated closely with the amount of IFN- γ mRNA. Abundant transcription of both macrophage products was a feature of granulomatous synovitis. Moderate levels were detected in follicular synovitis, and tissues with diffuse synovitis produced low concentrations of IL-1 β and TNF- α . The subdued production of the T cell products, IFN- γ and IL-4, as well as the macrophage products, IL-1 β and TNF- α , in samples with diffuse synovitis could not be explained by the action of the anti-inflammatory cytokines, IL-10 and TGF- β 1. High transcription of the IL-10 gene was typical for follicular disease. TCF- β 1 was most abundant in tissue extracts from follicular synovitis, but it was also present in high amounts in synovial biopsies with granuloma formation.

Taken together, reproducible patterns of organization of inflammatory cells in the synovial infiltrates were distinguished. The topography of the infiltrates correlated with distinct profiles of tissue cytokines, indicating that RA is not always characterized by a predominant T_H1 response as previously reported (Simon AK, *et al.*, *Proc Natl Acad Sci USA* 91:8562 (1994)) but that different cytokine pathways underlie the synovial inflammation (Table 7). RA is not a consequence of an immunodeviation of the cellular cytokine response as has been postulated for other autoimmune diseases (Finkelman FD, *J Exp Med* 182:279

(1995)), but it includes different response patterns, each of which may be genetically determined and may have different clinical implications.

Support for clinical relevance of the described patterns in organization and functional commitment of the infiltrating cells came from an analysis of disease manifestations and outcome. The best predictor for diffuse synovitis was a lack of RF production, a disease type previously associated with the genotype HLA-DRB1*01 or characterized by the absence of disease-associated HLA-DRB1 alleles. Patients with this form of RA synovitis often only require non-steroidal anti-inflammatory drugs and infrequently receive more aggressive treatment. Granulomatous synovitis was encountered in patients with extra-articular spreading and nodule formation. Comparison among different joints from the same patient have shown that a pattern is characteristic for each patient and that the synovial lesions in different joints display identical organizations.

The phenotypic dissection of rheumatoid synovitis requires the reevaluation of the concept that has recently been proposed as the major mechanism leading to tissue destructive inflammation. Immune deviation has been defined as an aberrant commitment to an immune pathway in response to a given antigen. Elegant studies in model systems such as leishmania major infection have provided evidence that, dependent on the genetic background of the host, an immune response to a given antigen can take different directions, thereby determining whether the disease resolves or progresses (Heinzel FP, *et al.*, *J Exp Med* 169:59 (1989)). It has been suggested that patients with RA display a similar immune deviation with a bias towards T_H1 responses (Simon AK, *et al.*, *Proc Natl Acad Sci USA* 91:8562 (1994)). Consequently, it has been proposed to use IL-4 and IL-10 as novel therapeutic agents in the attempt to restore T_H2 responsiveness and suppress dominant T_H1 responses (Isomaki P, *et al.*, *Arthritis Rheum* 39:386 (1996); van Roon JA, *et al.*, *Arthritis Rheum* 39:829 (1996); and Walmsley M, *et al.*, *Arthritis Rheum* 39:495 (1996)). The data herein describing three different tissue cytokine profiles raise doubt that RA can be considered a single entity caused by overproduction of T_H1 cytokines. This model may apply to only a subset of individuals clinically classified as RA. The proposed treatment approaches with IL-4 and IL-10 illustrate the shortcomings of a disease model of RA that does not adequately consider phenotypic and genotypic variability. IL-4 and IL-10 might be beneficial in some but not all forms of the

disease. Ignoring the heterogeneity of the disease process and its clinical spectrum could represent a major limitation in treatment trials.

Table 7. Tissue cytokine patterns in subtypes of rheumatoid synovitis

Cytokine	Diffuse synovitis	Follicular synovitis	Granulomatous synovitis
IFN- γ	+	++	+++
IL-4	+	-	+++
IL-1 β	+	++	+++
TNF- α	+	++	+++
IL-10	+	+++	+
TGF- β 1	++	+++	+++

Example 10 Evaluating RA

RA can be subsetted into different phenotypes that correlate with the inheritance of distinct combinations of disease-risk genes (Figure 6). Combinations of HLA-DRB1 alleles and particular polymorphisms of HLA-DRB1 genes are useful in dissecting patients with rheumatoid organ disease, the most feared complication of RA. Contribution from a distinct genetic system to rheumatoid organ disease is suggested by the accumulation of unusual lymphocytes in patients suffering from complications of RA. CD4 T cells characterized by the deficiency of CD28, a major co-stimulatory molecule, expand to high frequencies in patients with extra-articular RA. CD4⁺/CD28^{null} T cells are functionally active, produce high amounts of IFN- γ and use a yet unknown alternate costimulatory pathway. These cells typically undergo clonal proliferation *in vivo* and infiltrate into the inflammatory lesions. Concordance for the expansion of this unusual lymphocyte subset in monozygotic twins lends support to a genetic control of their emergence. CD4⁺/CD28^{null} T cells accumulate in RA patients independent of HLA-DR polymorphisms. Therefore, genes regulating their generation might represent novel RA risk genes.

Additional RA risk genes can include genes involved in regulating cytokine pathways.

Analysis of tissue cytokine profiles in rheumatoid synovitis demonstrated three distinct patterns of microanatomical organization of the inflammatory infiltrates and, more importantly, a close correlation of lymphoid structures and cytokine networks. RA does not simply arise from immune deviation; at least three different cytokine pathways are involved in RA synovitis. Recognition of this heterogeneity is essential in trying to identify the molecular mechanisms that induce either diffuse, follicular, or granulomatous synovitis.

Accepting the model that more than one genotype can be associated with RA has enormous implications. Genomic searches trying to identify shared denominators among affected individuals rely heavily on the definition of disease. If individuals with too many different types of disease are compared, the chance of identifying disease risk genes is low. Similar considerations apply to studies trying to identify disease instigators. If disease processes are heterogeneous and patients with different variants of RA are studied, it might be impossible to identify a causative agent present in all or most of them. All of these approaches would have a much higher power if study cohorts could be identified that represented distinct disease entities. Focusing on a single variant of RA would enhance the chance of identifying molecular abnormalities, genetic risk factors, and potential infectious agents involved in RA pathogenesis. From a clinical point of view, the dissection of phenotypic/genotypic variants of RA is critical for further exploration of new and more selective therapeutic avenues. The therapeutic benefit of intervention might depend on the ability to target the appropriate patient subset.

The cytokine profile within synovial tissue, the patient's HLA genotype, and the patient's frequency of CD4⁺/CD28^{null} T cells was used to determine the patient's predisposition to develop severe disease. Subcutaneous nodularity was used as an indicator of disease severity. Tissue type was independent of HLA-DRB1 genotype and CD4⁺/CD28^{null} T cells counts were independent of HLA-DRB1 genotype (Table 8). Granulomatous disease predicts nodularity independent of HLA-DRB1 genotype and CD4⁺/CD28^{null} T cells counts. With respect to diffuse disease, CD4⁺/CD28^{null} T cell counts were a better predictor for nodular disease than a HLA-DRB1 homozygous positive genotype.

Table 8. Evaluating RA patients for disease severity.

Tissue cytokines			
IL-4	low	low	high
IL-10	low	high	low
IFN- γ	low	high	high
Corresponding histology	diffuse	follicular	granulomatous
Patients (number)	31	12	6
RA-associated HLA-DRBI *04 allele	20 (65%)	7 (58%)	4 (66%)
Two RA-associated HLA-DRB1 alleles (Double-dose)	8 (25%)	3 (25%)	0 (0%)
Nodularity (%)	9 (29%)	2 (17%)	5 (84%)
% of nodular patient being double-dose (sensitivity)	2 (22%)	1 (50%)	0%
% of double-dose patients being nodular	2 (25%)	1 (33%)	N/A
CD4+ CD28- data available	7	3	3
Number of patients having high CD4 ⁺ /CD28 ^{null} counts	3	2	2
Number of patients being nodular	2	1	3
% of patients having high CD4 ⁺ /CD28 ^{null} counts that are nodular	2/3 (66%)	1 (50%)	100%

[illegible]

5 Other aspects, advantages, and modifications are within the scope of the following claims.

[illegible]

- 46

[illegible]

11. The method of claim 5, wherein said population comprises a population of patients having extra-articular involvement.

13. A method for determining the predisposition of a rheumatoid arthritis patient to develop severe disease, said method comprising:

- a) determining the level of a cytokine within a sample from said patient,
- b) determining the frequency of CD4⁺/CD28^{null} cells in said patient,
- c) comparing said level of said cytokine to a reference level and said frequency of CD4⁺/CD28^{null} cells to a reference frequency to obtain information about said predisposition, and
- d) determining if said patient is predisposed to develop severe disease based on said information.

15. The method of claim 13, wherein said sample comprises a tissue biopsy.

16. The method of claim 15, wherein said tissue biopsy comprises a synovial tissue biopsy.

17. The method of claim 13, wherein said reference level comprises the median level of

cytokine found in tissue samples derived from a population.

18. The method of claim 17, wherein said population comprises a population of patients having a diffuse rheumatoid arthritis condition.

19. The method of claim 17, wherein said population comprises a population of patients having a follicular rheumatoid arthritis condition.

20. The method of claim 17, wherein said population comprises a population of patients having a granulomatous rheumatoid arthritis condition.

21. The method of claim 17, wherein said population comprises a population of healthy individuals.

22. The method of claim 17, wherein said population comprises a population of patients having subcutaneous nodules.

23. The method of claim 17, wherein said population comprises a population of patients having extra-articular involvement.

24. The method of claim 17, wherein said population comprises a population of patients having major joint destruction.

25. The method of claim 13, wherein said frequency of $CD4^+/CD28^{null}$ cells comprises the percent of $CD4^+$ cells that are CD28 negative.

26. The method of claim 25, wherein said reference frequency is derived from the $CD4^+/CD28^{null}$ cell frequency from a population.

27. A method for determining the predisposition of a rheumatoid arthritis patient to

[illegible]

28. The method of claim 27, wherein said cytokine comprises a cytokine selected from the group consisting of IL-4, IL-10, and IFN- γ .

30. The method of claim 27, wherein said tissue biopsy comprises a synovial tissue biopsy.

32. The method of claim 31, wherein said population comprises a population of patients having a diffuse rheumatoid arthritis condition.

34. The method of claim 31, wherein said population comprises a population of patients having a granulomatous rheumatoid arthritis condition.

49

individuals.

36. The method of claim 31, wherein said population comprises a population of patients having subcutaneous nodules.

37. The method of claim 31, wherein said population comprises a population of patients having extra-articular involvement.

38. The method of claim 31, wherein said population comprises a population of patients having major joint destruction.

39. The method of claim 27, wherein said polymorphism comprises an HLA-DRB1 allele that encodes a polypeptide having an uncharged amino acid at position 74.

40. The method of claim 39, wherein said polymorphism comprises an HLA-DRB1 allele that encodes a polypeptide free from negatively charged amino acids at positions 70 and 71.

41. A method for determining the predisposition of a rheumatoid arthritis patient to develop severe disease, said method comprising:

- a) determining the level of a cytokine within a sample from said patient,
- b) determining the frequency of CD4⁺/CD28^{null} cells in said patient,
- c) comparing said level of said cytokine to a reference level and said frequency of CD4⁺/CD28^{null} cells to a reference frequency to obtain information about said rheumatoid arthritis condition,
- d) determining the presence of a polymorphism in an HLA-DRB1 allele in said patient, and
- e) determining if said patient is predisposed to develop severe disease based on said information and said presence of said polymorphism.

42. A kit for providing diagnostic information about a rheumatoid arthritis condition in a

patient, said kit comprising:

a) a binding pair member, wherein said binding pair member has specific binding affinity for a cytokine such that the level of said cytokine within a sample from said patient is determinable, and

b) a reference chart, wherein said reference chart contains information about cytokine levels such that an indication of the diffuse, follicular, or granulomatous nature of said rheumatoid arthritis condition is determinable based on said level of said cytokine within said sample.

43. A kit for providing diagnostic information about a rheumatoid arthritis condition in a patient, said kit comprising:

a) a binding pair member, wherein said binding pair member has specific binding affinity for a nucleic acid sequence encoding a cytokine such that the level of said cytokine within a sample from said patient is determinable, and

b) a reference chart, wherein said reference chart contains information about cytokine levels such that an indication of the diffuse, follicular, or granulomatous nature of said rheumatoid arthritis condition is determinable based on said level of said cytokine within said sample.

44. A kit for determining the predisposition of a rheumatoid arthritis patient to develop severe disease, said kit comprising:

a) a first binding pair member, wherein said first binding pair member has specific binding affinity for a cytokine or nucleic acid encoding said cytokine such that the level of said cytokine within a sample from said patient is determinable,

b) second binding pair member, wherein said second binding pair member has specific binding affinity for a $CD4^+/CD28^{null}$ cell marker such that the frequency of $CD4^+/CD28^{null}$ cells in said patient is determinable, and

c) a reference chart, wherein said reference chart contains information about cytokine levels and $CD4^+/CD28^{null}$ cell frequencies such that an indication of said predisposition is determinable based on said level of said cytokine within said sample and said frequency of

[illegible]

a) a binding pair member, wherein said binding pair member has specific binding affinity for a cytokine or nucleic acid encoding said cytokine such that the level of said cytokine within a sample from said patient is determinable,

c) a reference chart, wherein said reference chart contains information about cytokine levels such that an indication of said predisposition is determinable based on said level of said cytokine within said sample and said polymorphism of said patient.

47. A kit for determining the predisposition of a rheumatoid arthritis patient to develop severe disease, said kit comprising:

b) a second binding pair member, wherein said second binding pair member has specific binding affinity for a CD4⁺/CD28^{null} cell marker such that the frequency of CD4⁺/CD28^{null} cells in said patient is determinable,

d) a reference chart, wherein said reference chart contains information about cytokine levels and CD4⁺/CD28^{null} cell frequencies such that an indication of said predisposition is

determinable based on said level of said cytokine within said sample, said frequency of CD4⁺/CD28^{null} cells in said patient, and said polymorphism of said HLA-DRB1 allele.

ABSTRACT


The invention provides methods and materials for diagnosing a rheumatoid arthritis condition in a patient. Specifically, the invention provides methods and materials for classifying a rheumatoid arthritis condition as diffuse, follicular, or granulomatous. In addition, the invention provides methods and materials for determining if an individual suffering from a rheumatoid arthritis condition will develop severe disease.

24566.m11

FIGURE 1

IFN- γ gene expression

Cytic transcripts / 2×10^6 β -actin r



IL-4




FIGURE 2

IL-1 β and TNF- α







FIGURE 3


IL-10

TGF- β 1


DIFFUSE GRANULOMATOUS



0.000



... ..



<u>Country</u>	<u>Application No.</u>	<u>Filing Date</u>	<u>Priority Claimed</u>
----------------	------------------------	--------------------	-------------------------

Combined Declaration and Power of Attorney

Page 2 of 2 Pages

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Mark S. Ellinger,
Reg. No. 34,812
Ronald C. Lundquist,
Reg. No. 37,875
Rene D. Tegtmeyer,
Reg. No. 33,567

Dorothy P. Whelan,
Reg. No. 33,814
Monica McCormick Graham,
Reg. No. 42,600
Eldora L. Ellison,
Reg. No. 39,967

Richard J. Anderson,
Reg. No. 36,732
J. Patrick Finn III,
Reg. No. 44,109

Address all telephone calls to Mark S. Ellinger at telephone number 612-335-5070.

Address all correspondence to Mark S. Ellinger at:

FISH & RICHARDSON P.C., P.A.
60 South Sixth Street, Suite 3300
Minneapolis, Minnesota 55402

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

Full Name of Inventor: Jorg J. Goronzy

Inventor's Signature:

Residence

Address:

Citizenship:

Post Office

Address:

1804 Spel Lane S.W.

Rochester, Minnesota 55902

German

200 First Street S.W.

Rochester, Minnesota 55905

Date: Nov. 5, 99

Full Name of Inventor: Cornelia M. Weyand

Inventor's Signature:

Residence

Address:

Citizenship:

Post Office

Address:

1804 Spel Lane S.W.

Rochester, Minnesota 55902

German

200 First Street S.W.

Rochester, Minnesota 55905

Date: Nov 5, 99